

**OVERT AGITATION SEVERITY SCALE (OASS)**  
Yudofsky SC ,Kopecky HJ,Kunik M,Silver JM,Endicott J

INTENSITY (I)	Behavior	Frequency(f)					Severity score(ss)(IxF=ss)
A.	Vocalization & Oral / Facial Movements	NOT PRESENT	RARELY	SOME OF THE TIME	MOST OF THE TIME	ALWAYS PRESENT	
1	Whispering, Whining, Moaning, grunting, Crying	0	1	2	3	4	0
2	Smacking or Licking of lips, Chewing, ClenchingJaw, Licking, grimacing, apitting	0	1	2	3	4	0
3	Rocking, Twisting, banging of Head	0	1	2	3	4	0
4	Vocal Perseverating, Screaming, Cursing, Threating, Wailing	0	1	2	3	4	0
B.	Upper Torso & Upper Extremity Movements						
1	Tapping fingers, fideting, or wringing of hands, Swinging or flailing arms	0	1	2	3	4	0
2	Teak Persevarating(e.g , opening and Closing drawers, folding and unfolding clothes, picking at objects, clothes or self, pulling at own hair)	0	1	2	3	4	0
3	Rocking (back & forth), bobbing(up and down)twisting, writhing of torso, rubbing or masturbating self	0	1	2	3	4	0
4	Slapping, Swatting, hitting at objects or others	0	1	2	3	4	0
C.	Lower Extremity Movements						
1	Trapping toes, Clenchig toes, Tapping Heels, Extending, Flaxing or Twisting Foot	0	1	2	3	4	0
2	Shaking Legs, tapping Knees and/or things, thrusting Pelvis, stomping	0	1	2	3	4	0
3	Pacing, wandoring	0	1	2	3	4	0
4	Thrashing legs, Kicking at objects or others	0	1	2	3	4	0

Total Gass	[ ]
Subtract Base lineGass	[ ]
Revised Gass	[ ]

**Instruction for Completing Form**

- Step One: For each behaviour,circle the corresponding frequency after 15 minutes of observation.
- Step Two: For every behavior exhibited multiply the Intensity score (I) by the Frequency (f) and record as the severity Score (SS).
- Step Three: For the OVERT AGITATION SEVERITY SCORE (OASS) total all severity scores and record as Total OASS.
- Step Four: Does thia patient have a Neuromuscular Diaorder (i.e., Parkinson's Disease, tardive dyskinesia) affecting Total OASS? Yes No
- Step Five: If yes, please establish a baselina OASS in non-agitated state and subtract from above Total 0ASS for Revised OASS. Total 0ASS for Revised OASS.

**COMMENTS:**

katana

DIAGNOSIS: F25.0 - SCHIZOAFFECTIVE DISORDER, BIPOLAR TYPE	NAME OF RATER: Reed, Jody
SEX OF PATIENT: Male	TIME OF OBSERVATION: 15:06:17
AGE: 30	DATE: 11/05/2021

CURRENT MEDICATION:		
Name: Ativan	Dose: 2 mg	Frequency: PRN Q4HOUR
Name: Seroquel	Dose: 100 mg	Frequency: BID